



Orbital Tori Construction Using Trajectory Following Spectral Methods

By Ralph E. Bordner

Biblioscholar Okt 2012, 2012. Taschenbuch. Book Condition: Neu. 246x189x11 mm. This item is printed on demand - Print on Demand Neuware - The Kolmogorov-Arnold-Moser (KAM) theorem states that lightly perturbed integrable Hamiltonian systems maintain their multiply periodic, toroidal motion in the phase space. The assertion that earth orbiting satellites under the influence of the geopotential mimic this behavior is the underlying premise of this work. This paper focuses on applying trajectory following spectral methods on selected orbits to decompose them into multi-periodic Fourier series, effectively compressing ephemerides for long-term use. The proposed approach focuses on fitting local spectral structures, denoted as frequency clusters, within the sampled orbital data to the analytical form of the windowed, truncated, continuous Fourier transform. This approach is significantly more numerically efficient than fitting every coefficient within the N-tuple Fourier series simultaneously. Numerical results using integrated data show maximum error in orbital torus fits can be kept at the single-digit-meter level per coordinate axis over a 1 year period. 188 pp. Englisch.



READ ONLINE
[1.34 MB]

Reviews

A top quality ebook and the typeface used was interesting to read through. It is rally intriguing throug reading through period. You wont feel monotonny at anytime of the time (that's what catalogues are for relating to when you ask me).

-- Estelle Donnelly

Without doubt, this is the best operate by any publisher. I was able to comprehended everything out of this written e publication. Its been developed in an remarkably easy way which is only following i finished reading through this ebook by which basically altered me, modify the way i believe.

-- Dr. Ofelia Grant Sr.